



PROPOZYCJA
PLANU AGLOMERACJI MIASTA PRZEMYSŁA
Aktualizacja 2013



-  Pencil icon (drawing)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)
 Pencil icon (eraser)

Ochsen

[illegible]

WT: 25000.4 576.32
Proteomyl3



5 9 0 0 5 7 5 0 1 8 9 0 1

1 : 25 000

Figure 1 is a line graph showing the change in the number of individuals of the 1st and 2nd instars of the European spruce sawfly (*Pristiphora abietis*) over time. The x-axis represents time in days (0 to 1000). The y-axis represents the number of individuals in thousands (0 to 1000). Two lines are plotted: a solid line for the 1st instar and a dashed line for the 2nd instar. The 1st instar population peaks around day 200 and then declines. The 2nd instar population peaks around day 400 and then declines.

Opisano u CPSE Knjizi 1987:
na području mapy 1:10 000 u 1983;
"na stadiju izmjereno" - u 1989;
Dok
Wypasa jezera